

Note to readers with disabilities: *EHP* strives to ensure that all journal content is accessible to all readers. However, some figures and Supplemental Material published in *EHP* articles may not conform to [508 standards](#) due to the complexity of the information being presented. If you need assistance accessing journal content, please contact ehp508@niehs.nih.gov. Our staff will work with you to assess and meet your accessibility needs within 3 working days.

Supplemental Material

Association of Cardiometabolic Genes with Arsenic Metabolism Biomarkers in American Indian Communities: The Strong Heart Family Study (SHFS)

Poojitha Balakrishnan, Dhananjay Vaidya, Nora Franceschini, V. Saroja Voruganti, Matthew O. Gribble, Karin Haack, Sandra Laston, Jason G. Umans, Kevin A. Francesconi, Walter Goessler, Kari E. North, Elisa Lee, Joseph Yracheta, Lyle G. Best, Jean W. MacCluer, Jack Kent, Jr, Shelley A. Cole, and Ana Navas-Acien

Table of Contents

- Table S1.** Inter-batch variability of urine arsenic measurements (n=46)
- Table S2.** Fine mapped loci on MetaboChip
- Table S3.** Top MetaboChip associations for percent inorganic arsenic
- Table S4.** Top MetaboChip associations for percent monomethylarsonate
- Table S5.** Top MetaboChip associations for percent dimethylarsinate
- Table S6.** Top MetaboChip associations for principal component 1 of arsenic species
- Table S7.** Top MetaboChip associations for principal component 2 of arsenic species
- Table S8.** Top candidate SNP associations for percent inorganic arsenic
- Table S9.** Top candidate SNP associations for percent monomethylarsonate
- Table S10.** Top candidate SNP associations for percent dimethylarsinate
- Table S11.** Top candidate SNP associations for principal component 1 of arsenic species

Table S12. Top candidate SNP associations for principal component 2 of arsenic species

Figure S1. Quantile-quantile plot for MetaboChip of percent arsenic species and principal components of arsenic species

Figure S2. Quantile-quantile plot for candidate SNPs of percent arsenic species and principal components of arsenic species

Figure S3. Manhattan plot of MetaboChip associations for percent inorganic arsenic

Figure S4. Manhattan plot of MetaboChip associations for percent monomethylarsononate

Figure S5. Manhattan plot of MetaboChip associations for percent dimethylarsinate

Figure S6. Manhattan plot of MetaboChip associations for principal component 1 of arsenic species

Figure S7. Manhattan plot of MetaboChip associations for principal component 2 of arsenic species

Figure S8. Regional association plot at 10q24 of percent arsenic species conditioned on rs12768205

Figure S9. Regional association plot at 10q24 of principal components of arsenic species conditioned on index SNP

Figure S10. Distribution of percent arsenic species by rs3740394 genotype

Figure S11. Distribution of percent arsenic species by rs7098825 genotype